

EFFECTIVE ACTION FOR DIRAC SPINORS IN THE UNIFORM ELECTROMAGNETIC BACKGROUND FIELDS

ROBERTO SOLDATI

Dipartimento di Fisica "A. Righi", Università di Bologna,

LORENZO SORBO

S.I.S.S.A.-I.S.A.S., Trieste

Some new expressions are found, concerning the effective action as a regularized path-integral for Dirac's spinors in $3+1$ dimensions, in the presence of general uniform (i.e., constant and homogeneous) electric and magnetic fields. The rate of e^+e^- pairs production is computed and briefly discussed.

Path-integrals over Grassmann-valued Fermi fields are formally given by means of the well known Berezin's recipe¹. Gaussian path-integrals of the above kind involving some self-adjoint Dirac-like operator are *naïvely* equivalent to functional determinants; for instance

$$\int [d\bar{\psi}][d\psi] \exp \left\{ - \int \bar{\psi} D \psi dx \right\} = \prod_n \lambda_n = \det D, \quad (1)$$

where $D = D^\dagger$ and $D\psi_n = \lambda_n\psi_n$, $\lambda_n \neq 0$, $\{\psi_n\}$ being the complete orthonormal set of eigenfunctions of some self-adjoint invertible operator D with purely discrete non-degenerate spectrum.

Functional determinants can be defined in a rigorous mathematical way² for general dimensionless invertible elliptic systems of partial differential operators defined on some compact manifold without boundary. For such^a a system of operators A :

(i) **complex** powers A^{-s} do exist, (ii) A^{-s} has a **continuous kernel** K_{-s} , if the real part of s is sufficiently large, (iii) $K_{-s}(x, y)$ is an **entire** function of s , where (x, y) are local coordinates on a compact manifold $\mathcal{M}$ without boundary and $x \neq y$, (iv) $K_{-s}(x, x)$ extends to a **meromorphic** function, (v) $K_{-s}(x, x)$ has **no poles** at $s = 0, -1, -2, \dots$

Owing to the above results, we can safely define the ζ -function associated to the system A as

$$\zeta_A(s) \equiv \int_{\mathcal{M}} \text{tr} K_{-s}(x, x) d\mu(x), \quad (2)$$

^aThe system A is supposed to be elliptic of order $m > 0$, invertible and to admit a ray of minimal growth for its resolvent.

where the trace tr refers to discrete matrix-like indices of the system A while $d\mu$ is the invariant measure on the compact manifold $\mathcal{M}$. Thanks to (v), the functional determinant of A can be rigorously defined³ by means of

$$\mathbf{det} A \equiv \exp \left\{ -\frac{d}{ds} \zeta_A(s=0) \right\}. \quad (3)$$

Let us now consider quantized Dirac's four components massive spinor fields in the presence of some background classical uniform (i.e., constant and homogeneous) general electromagnetic field. The use of the path-integral method forces us to work in the euclidean framework: only at the very end of our calculations we shall operate a Wick rotation to Minkowski space-time^b. The effective euclidean action in the semi-classical approximation is given by

$$\mathcal{S}_{\text{Eff}}^{\text{E}}[A_\mu] = \int d^4x \frac{1}{4} F_{\mu\nu} F_{\mu\nu} - \log \mathbf{det} \left(\frac{D[A_\mu] + im}{\mu} \right), \quad (4)$$

where $D[A_\mu] \equiv \gamma_\mu (i\partial_\mu + eA_\mu)$ is the self-adjoint euclidean Dirac operator, m being the fermion mass and μ some suitable mass scale. As the massive Dirac's operator is normal we have

$$\left| \mathbf{det} \left(\frac{D + im}{\mu} \right) \right|^2 = \mathbf{det} \left(\frac{D^2 + m^2}{\mu^2} \right) \equiv \mathbf{det} H. \quad (5)$$

The effective lagrangean is thereby defined to be

$$\mathcal{L}_{\text{Eff}}^{\text{E}}[A_\mu] = \mathcal{L}_{\text{Cl}}^{\text{E}}[A_\mu] - \frac{1}{2} \mu^4 \left. \frac{\partial}{\partial s} \zeta_H(s) \right|_{s=0}. \quad (6)$$

Since the square H of the massive Dirac's operator is elliptic, invertible and positive, its determinant can be evaluated by means of the ζ -regularization. A frame can always be chosen such that $F_{03} = -F_{30} = E$, $F_{12} = -F_{21} = B$, all the other components of $F_{\mu\nu}$ vanishing.

After setting $a \equiv |eE/\mu^2|$, $b \equiv |eB/\mu^2|$ and $c \equiv (m^2/\mu^2)$ we get

$$\zeta_H(s) = \frac{ab}{4\pi^2 \Gamma(s)} \int_0^\infty dt t^{s-1} e^{-ct} \left\{ 2 \frac{e^{-2at} + e^{-2bt}}{(1 - e^{-2at})(1 - e^{-2bt})} + 1 \right\}. \quad (7)$$

Explicit formulae were known⁴ for the effective euclidean lagrangean in the cases $B = 0$ (or $E = 0$) and $E = B$. It is possible to obtain asymptotic expansions⁵ in powers of $|B/E|$ both in the massive and massless cases.

^bOn $\mathbf{R}^4$ one can rigorously define the effective action per unit volume.

The starting point is the scaling property $\zeta_H(s) = a^{2-s}g(s; y, z)$, so that it is sufficient to study the relevant quantity

$$g(s; y, z) = \frac{y}{\Gamma(s)} \int_0^\infty dt t^{s-1} e^{-zt} \left\{ 2 \frac{e^{-2t} + e^{-2yt}}{(1 - e^{-2t})(1 - e^{-2yt})} + 1 \right\}, \quad (8)$$

as a function of the ratio $y \equiv (b/a) = |B/E|$ at a given $z \equiv (c/a) = |m^2/eE|$. Expansions around $y = 1 \Leftrightarrow |E| = |B|$ are easily obtained. The non-trivial result is the expansion around $|B| = y = 0$:

$$g(s; y, z) = \frac{2^{1-s}}{\Gamma(s)} \sum_{k=0}^N \frac{\mathbf{B}_{2k}}{(2k)!} y^{2k} \Gamma(s-1+2k) \left\{ 2\zeta_R(s-1+2k; 1 + \frac{z}{2}) + \left(\frac{z}{2}\right)^{1-2k-s} \right\} + O(y^{2N+2}), \quad (9)$$

in which $\mathbf{B}_{2k}$ is the $2k$ -th Bernoulli number and $\zeta_R(a; x)$ is the Riemann-Hurwitz ζ -function⁶. Exploiting the above scaling property the effective euclidean lagrangean of QED in the presence of uniform fields can be easily calculated as an asymptotic series in $|B/E|$ or $|E/B|$.

Now we show how these corrections to the $E = 0$ case work, paying special attention to the rate of production of fermion-antifermion pairs. Denoting by $\mathcal{E}$ and $\mathcal{B}$ the minkowskian electric and magnetic field strengths respectively, the transition to the Minkowski space-time is performed by means of the substitution

$$\mathcal{L}_{\text{Eff}}^{\text{M}}(\mathcal{E}, \mathcal{B}) = -\mathcal{L}_{\text{Eff}}^{\text{E}}(E = -i\mathcal{E}, B = \mathcal{B}). \quad (10)$$

The exact pairs production rate per unit volume $w = 2Im\mathcal{L}_{\text{Eff}}^{\text{M}}$ at $\mathcal{B} = 0$ is⁷

$$w(\mathcal{E}, \mathcal{B} = 0) = \frac{e^2 \mathcal{E}^2}{4\pi^3} \sum_{n=1}^{\infty} \frac{1}{n^2} \exp \left\{ -n \frac{\pi m^2}{e\mathcal{E}} \right\}, \quad (11)$$

which shows that the function $w(\mathcal{E})$ is not analytic in $\mathcal{E} = 0$. Now, with the aid of the previous asymptotic expansion we obtain the first corrections of the order $O(\mathcal{B}/\mathcal{E})^2$ to the total rate of production of e^+e^- pairs that reads

$$w = \frac{e^2 \mathcal{E}^2}{4\pi^3} \sum_{n=1}^{\infty} \left(\frac{1}{n^2} + \frac{\pi^2}{3} \frac{\mathcal{B}^2}{\mathcal{E}^2} \right) \exp \left\{ -n \frac{\pi m^2}{e\mathcal{E}} \right\} + O\left(\frac{\mathcal{B}^4}{\mathcal{E}^4}\right). \quad (12)$$

The above formulae are not analytic in $\mathcal{E} = 0$, owing to the non-perturbative character of this phenomenon. Considering also the second term of order $O(\mathcal{B}/\mathcal{E})^4$ to the "unperturbed" effective lagrangean and retaining only the

first terms in the expansion for weak fields, we get the effective lagrangean in the limit $|e\mathcal{B}| \ll |e\mathcal{E}| \ll m^2/2$: namely,

$$\delta^{(4)}\mathcal{L}_{\text{Eff}}^{\text{M}}(\mathcal{E}, \mathcal{B}) \simeq \frac{e^4}{16\pi^2} \frac{2}{45m^4} (\mathcal{E}^4 + \mathcal{B}^4 + 5\mathcal{E}^2\mathcal{B}^2) + \frac{e^2}{24\pi^2} \mathcal{B}^2 \log \frac{m^2}{\mu^2}, \quad (13)$$

which is in perfect agreement with the Euler-Heisenberg effective lagrangean⁸ describing the light-light scattering at very low photons momenta, after setting the arbitrary scale μ at the main scale m of the problem. According to a *naïve* interpretation, the phenomenon of particle production in external electromagnetic fields is due to the fact that virtual pairs created in the vacuum are accelerated by the electric field and may gain energy enough to reach the threshold (the electron/positron mass) and become physical particles. On the other hand electrically charged particles do not acquire energy from a magnetic field, so that it is rather surprising that a (weak) magnetic field gives itself a contribution to the rate of pair creation as we have shown to be

$$(\Delta w)_{\text{magn}} = \frac{e^2\mathcal{B}^2}{12\pi} \sum_{n=1}^{\infty} \exp \left\{ -n \frac{\pi m^2}{e\mathcal{E}} \right\} + O \left(\frac{\mathcal{B}^4}{\mathcal{E}^4} \right). \quad (14)$$

Analogous calculations in the non-Abelian case as well as in the case of chiral fermions would be quite suggestive and useful.

References

1. F. A. Berezin : " *The method of second quantization* ", Academic Press, New York (1966).
2. R. T. Seeley : Amer. Math. Soc. Proc. Symp. Pure Math. **10** (1967) 288.
3. D. B. Ray, I. M. Singer : Adv. Math. **7** (1971) 145; J. S. Dowker, R. Critchley : Phys. Rev. D **13** (1976) 3224; S. W. Hawking : Comm. Math. Phys. **55** (1977) 133.
4. S. K. Blau, M. Visser, A. Wipf : Int. Jour. Mod. Phys. A **6** (1991) 5409.
5. R. Soldati, L. Sorbo : Phys. Lett. B **426** (1998) 82.
6. A. Erdélyi, W. Magnus, F. Oberhettinger, F. G. Tricomi (Eds.) : " *The Bateman manuscript project: higher transcendental functions* ", McGraw-Hill, New York (1953-1955).
7. C. Itzykson, J. B. Zuber : " *Quantum field theory* ", McGraw-Hill, New York (1980).
8. H. Euler, W. Heisenberg : Z. Phys. **98** (1936) 714.